

Covering cubes by hyperplanes

Hao Huang

Emory University

Oct 8, 2020

Joint work with Alexander Clifton (Emory).



Alexander Clifton
(Ph.D student at Emory)

A naive question

The n -dimensional **cube** Q^n consists of the binary vectors $\{0, 1\}^n$.

An **affine hyperplane** is:

$$\{\vec{x} : a_1x_1 + \cdots + a_nx_n = b\}.$$

QUESTION

What is the minimum number of affine hyperplanes that cover all the vertices of Q^n ?

A naive question

The n -dimensional **cube** Q^n consists of the binary vectors $\{0,1\}^n$.


An **affine hyperplane** is:

$$\{\vec{x} : a_1x_1 + \dots + a_nx_n = b\}.$$

QUESTION

What is the minimum number of affine hyperplanes that cover all the vertices of Q^n ?

Answer: 2.



$$x_i = 1$$


$$x_i = 0$$

The Alon-Füredi Theorem

A NEW QUESTION

Suppose we would like to avoid exactly one vertex of the cube, how many affine hyperplanes are needed?

For Q^3 , 3 planes are needed.



THEOREM (ALON, FÜREDI 1993)


Any set of affine hyperplanes that covers all the vertices of the n -cube Q^n but one contains at least n affine hyperplanes.

The Alon-Füredi Theorem

A NEW QUESTION

Suppose we would like to avoid exactly one vertex of the cube, how many affine hyperplanes are needed?

For Q^3 , 3 planes are needed.



THEOREM (ALON, FÜREDI 1993)


Any set of affine hyperplanes that covers all the vertices of the n -cube Q^n but one contains at least n affine hyperplanes.

The Alon-Füredi Theorem

A NEW QUESTION

Suppose we would like to avoid exactly one vertex of the cube, how many affine hyperplanes are needed?

For Q^3 , 3 planes are needed.



THEOREM (ALON, FÜREDI 1993)


Any set of affine hyperplanes that covers all the vertices of the n -cube Q^n but one contains at least n affine hyperplanes.

The Alon-Füredi Theorem

A NEW QUESTION

Suppose we would like to avoid exactly one vertex of the cube, how many affine hyperplanes are needed?

For Q^3 , 3 planes are needed.



THEOREM (ALON, FÜREDI 1993)


Any set of affine hyperplanes that covers all the vertices of the n -cube Q^n but one contains at least n affine hyperplanes.

The Alon-Füredi Theorem

A NEW QUESTION

Suppose we would like to avoid exactly one vertex of the cube, how many affine hyperplanes are needed?

For Q^3 , 3 planes are needed.



THEOREM (ALON, FÜREDI 1993)

Any set of affine hyperplanes that covers all the vertices of the n -cube Q^n but one contains at least n affine hyperplanes.

An outline of the proof of Alon-Füredi Theorem

Suppose

Proof. H_1, \dots, H_m affine hyperplanes covers $\mathbb{Q}^n \setminus \{0\}$

$$0 \notin H_i$$

$$H_i: \langle \vec{x}, \vec{a}_i \rangle = b_i \quad b_i \neq 0$$

$\vec{x} \in \mathbb{R}^n \quad \vec{a}_i \in \mathbb{R}^n$

$$p(x_1, \dots, x_n) = \prod_{i=1}^m (\langle \vec{x}, \vec{a}_i \rangle - b_i)$$

① $p(\vec{x}) = 0$ for $\vec{x} \in \mathbb{Q}^n \setminus \{0\}$

② $p(\vec{0}) \neq 0$

$\deg p = m$

p is in n vars.

$$\Rightarrow m \geq n$$

$p \neq 0$ at $\vec{0}$
 $= 0 \quad \mathbb{Q}^n \setminus \{0\}$

$$p = (x_1 - 1) \cdots (x_n - 1)$$

$$x_1 = 1$$

$$\vdots$$

$$x_{n-1} = 1$$

$$x_1 + \dots + x_n = 1$$

p

Covering the cube twice

QUESTION (BUKH'S HOMEWORK ASSIGNMENT AT CMU)

What happens if we would like to cover the vertices of Q^n at least **twice**, with one vertex uncovered?

$2n$ $x_1=1$
 \vdots twice.
 $x_n=1$

$\geq n+1$

Remove **one** hyperplane.

$Q^n - \{0\} \geq 1.$

$\frac{n+1}{e_i}$ $\left(\begin{matrix} x_1=1 \\ \vdots \\ x_n=1 \end{matrix} \right)$ $(, \dots, 1, \dots, 1, \dots)$
 \uparrow \uparrow
 i -th j -th
 $x_i=1$
 $x_j=1$
 $x_1 + x_2 + \dots + x_n = 1$

$\geq n$

$\geq n+1$

Covering the cube k times

Denote by $f(n, k)$ the minimum number of affine hyperplanes needed to cover every vertex of Q^n at least k times (except for $\vec{0}$ which is not covered at all).

We call such a cover an **almost k -cover** of the n -cube.

$$f(n, 1) = n.$$

$$f(n, 2) = n + 1.$$

What is the next?

Upper and lower bounds

$$f(n, k) \leq n + \binom{k}{2}$$

Take

$$\begin{array}{c}
 n \\
 + k-1 \\
 + k-2 \\
 + \vdots \\
 + 1 \\
 \hline
 \binom{k}{2} + n
 \end{array}$$

$$x_1 = 1, \dots, x_n = 1,$$

$$x_1 + \dots + x_n = 1 \text{ for } k-1 \text{ times,}$$

$$x_1 + \dots + x_n = k-2 \text{ for twice.}$$

$$x_1 + \dots + x_n = k-1 \text{ for 1 time.}$$

If \vec{v} has t coordinates $= 1$.

then \vec{v} has already been counted t times

$$f(n, k) \geq n + k - 1$$

Note that removing $k-1$ planes from an almost k -cover still gives an almost 1-cover.

$$k = 3: \quad n + 2 \leq f(n, 3) \leq n + 3.$$

The $k = 3$ case and a natural conjecture

$$\begin{array}{ll} \pi_i = 1 & \text{twice} \\ \sum \pi_i = 1 & \text{once} \\ \sum \pi_i = 2 & \end{array}$$

THEOREM (H., CLIFTON 2019)

For $n \geq 2$,

$$f(n, 3) = n + 3.$$

$$n+k-1 = n+3$$

For $n \geq 3$,

$$f(n, 4) \in \{n + 5, n + 6\}.$$

$$n + \binom{k}{2} = n + 6$$

CONJECTURE (H., CLIFTON 2019)

For fixed integer $k \geq 1$ and sufficiently large n ,

$$f(n, k) = n + \binom{k}{2}.$$

$\geq k$ times

$$\boxed{kn} \leq$$

if n small

THE NULLSTELLENSATZ

If \mathbb{F} is an algebraically closed field, and $f, g_1, \dots, g_m \in \mathbb{F}[x_1, \dots, x_n]$, where f vanishes over all common zeros of g_1, \dots, g_m , then there exists an integer k , and polynomials $h_1, \dots, h_m \in \mathbb{F}[x_1, \dots, x_n]$, such that

$$f^k = \sum_{i=1}^m h_i g_i.$$

When $m = n$, and $g_i = \prod_{s \in S_i} (x_i - s)$, for some $S_1, \dots, S_n \subset \mathbb{F}$, a stronger result holds: there are polynomials h_1, \dots, h_n with $\deg h_i \leq \deg f - \deg g_i$, such that

$$f = \sum_{i=1}^n h_i g_i.$$

Punctured Combinatorial Nullstellensatz

We say $\vec{a} = (a_1, \dots, a_n)$ is a **zero of multiplicity t** of $f \in \mathbb{F}[x_1, \dots, x_n]$, if t is the minimum degree of the terms in $f(x_1 + a_1, \dots, x_n + a_n)$.

For $i = 1, \dots, n$, let

$$f = (x-1)(y-1) \quad \text{at } (1,1) \text{ it has multiplicity } 2$$

$$D_i \subset S_i \subset \mathbb{F}. \quad g_i = \prod_{s \in S_i} (x_i - s). \quad \ell_i = \prod_{d \in D_i} (x_i - d).$$

for $\{0,1\}$ $g_i = (x_i-1)x_i$ $\ell_i = x_i$

THEOREM (BALL, SERRA 2009)

If f has a zero of multiplicity at least t at all the common zeros of g_1, \dots, g_n , except at least one point of $D_1 \times \dots \times D_n$ where it has a zero of multiplicity less than t , then there are polynomials h_τ satisfying $\deg(h_\tau) \leq \deg(f) - \sum_{i \in \tau} \deg(g_i)$, and a non-zero polynomial u satisfying $\deg(u) \leq \deg(f) - \sum_{i=1}^n (\deg(g_i) - \deg(\ell_i))$, such that

$$f = \sum_{\tau \in T(n,t)} g_{\tau(1)} \cdots g_{\tau(t)} h_\tau + u \prod_{i=1}^n \frac{g_i}{\ell_i}.$$

$u \cdot \prod (x_i-1)$

$T(n, t)$ consists of all non-decreasing sequences of length t on $[n]$.

Outline of our proof using the PCN ($k = 3$)

Goal. $f(n, 3) \geq n+3$

Suppose not H_1, \dots, H_{n+2} $H_i: \langle \vec{x}, \vec{a}_i \rangle = 1$.

Let $p_i = \langle \vec{x}, \vec{a}_i \rangle - 1$. $f = p_1 \dots p_{n+2}$ ^{deg = n+2}

PCN \Rightarrow

$$f = \sum_{1 \leq i < j \leq n+1} x_i(x_i-1)x_j(x_j-1)x_{n+1}(x_{n+1}-1) + \left[\prod_{i=1}^n (x_i-1) \right] u$$

$$f = 0 \text{ on } \mathbb{Q}^n \setminus \{0\}$$

$$\frac{\partial f}{\partial x_i} \equiv \frac{\partial^2 f}{\partial x_i \partial x_j}$$


$$\Rightarrow u = 0$$

$$\deg u \leq \deg f - n$$

$$\leq 2$$

$$f(\vec{0}) \stackrel{(-1)^n}{=} u(\vec{0}) = 0$$

Follow-up work

$$x_1 = 1, \dots, x_{n-1} = 1 \quad x_1 + \dots + x_n = 1$$


The essence of this proof can be summarized in one sentence:

If f has zeroes of multiplicity at least 3 at $\{0, 1\}^n \setminus \{0\}$ and $f(0) \neq 0$, then $\deg(f) \geq n + 3$.

$$f = p_1 \cdots p_m$$

THEOREM (SAUERMAN, WIGDERSON 2020)

For $k \geq 2$, the minimum possible degree of a polynomial $f(x_1, \dots, x_n)$ such that it has zeroes of multiplicity at least k at $\{0, 1\}^n \setminus \{0\}$ and $f(0) \neq 0$, is $n + 2k - 3$.

$$n + \binom{k}{2} \geq f(n, k) \geq \underline{n + 2k - 3}$$

COROLLARY (SAUERMAN, WIGDERSON 2020)

For $k \geq 2$, an almost k -cover of Q^n has at least $n + 2k - 3$ hyperplanes.

$f(n, k)$ for fixed n and large k

For small n , $f(n, k) \neq n + \binom{k}{2}$. Actually,

THEOREM (H., CLIFTON 2019)


For fixed n , and k tends to infinity,

$$f(n, k) = \left(1 + \frac{1}{2} + \cdots + \frac{1}{n} + o(1)\right) k.$$

- Upper bound: use every hyperplane

$$x_{i_1} + \cdots + x_{i_j} = 1$$

a total of $\frac{k}{f_j(n)}$ times. e.g.




$$\frac{3k}{2}$$

$$\frac{3}{2} = 1 + \frac{1}{2}$$

$f(n, k)$ for fixed n and large k (ctd.)

- Lower bound: (e.g. $n = 3$) assign weights to vertices:



Every affine plane covers vertices of total weight at most 1.
Therefore one needs at least


$$k \cdot \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{3} \right) = \frac{11}{6}k$$

hyperplanes.

For general n , assign weight $1/(j \binom{n}{j})$ to vertices whose sum of coordinate is j .

$f(n, k)$ for fixed n and large k (ctd.)

- Lower bound: (e.g. $n = 3$) assign weights to vertices:



Every affine plane covers vertices of total weight at most 1.
Therefore one needs at least


$$k \cdot \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{3} \right) = \frac{11}{6}k$$

hyperplanes.

For general n , assign weight $1/(j \binom{n}{j})$ to vertices whose sum of coordinate is j .

$f(n, k)$ for fixed n and large k (ctd.)

- Lower bound: (e.g. $n = 3$) assign weights to vertices:



Every affine plane covers vertices of total weight at most 1.
Therefore one needs at least


$$k \cdot \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{3} \right) = \frac{11}{6}k$$

hyperplanes.

For general n , assign weight $1/(j \binom{n}{j})$ to vertices whose sum of coordinate is j .

$f(n, k)$ for fixed n and large k (ctd.)

- Lower bound: (e.g. $n = 3$) assign weights to vertices:



Every affine plane covers vertices of total weight at most 1.
Therefore one needs at least

$$k \cdot \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \frac{1}{6} + \frac{1}{6} + \frac{1}{6} + \frac{1}{3} \right) = \left(\frac{11}{6} \right) k = 1 + \frac{1}{2} + \frac{1}{3}.$$

hyperplanes.

For general n , assign weight $1/(j \binom{n}{j})$ to vertices whose sum of coordinate is j .

An LYM-like inequality

THE LUBELL-YAMATO-MESHALKIN INEQUALITY

Let \mathcal{F} be a family of subsets in which no set contains another, then

$$\sum_{S \in \mathcal{F}} \frac{1}{\binom{n}{|S|}} \leq 1.$$

LEMMA (H., CLIFTON 2019)

Given n real numbers a_1, \dots, a_n , let

if $a_i > 0$

$$\mathcal{F} = \left\{ S : \emptyset \neq S \subset [n], \sum_{i \in S} a_i = 1 \right\},$$

then

$$\sum_{S \in \mathcal{F}} \frac{1}{|S| \binom{n}{|S|}} \leq 1.$$

The inequality is tight for all non-zero binary (a_1, \dots, a_n) .

$$\frac{1}{j \binom{n}{j}}$$

$$(a_1 x_1 + \dots + a_n x_n) = 1$$

Proof of the Lemma

We associate every $S \in \mathcal{F}$ (binary vector covered by the plane) with some permutations in $\mathcal{P}_S \subset S_n$.

e.g. When $n = 5$, $S = \{1, 3, 4\}$, it means $a_1 + a_3 + a_4 = 1$, take all permutations in S_5 with prefix (i_1, i_2, i_3) satisfying

$$\{i_1, i_2, i_3\} = \{1, 3, 4\}, \quad a_{i_1} < 1, \quad a_{i_1} + a_{i_2} < 1.$$

We can show:

- \mathcal{P}_S are pairwise disjoint.
- $|\mathcal{P}_S| \geq (|S| - 1)!(n - |S|)!$ (the proof uses the *lorry driver puzzle*.)
- Therefore

$$n! \geq \sum_{S \in \mathcal{F}} |\mathcal{P}_S| = \sum_{S \in \mathcal{F}} (|S| - 1)!(n - |S|)!,$$

which simplifies to our desired result.

Future research problems (I)

$$n + \binom{k}{2}$$

$$n + 2k - 3$$

PROBLEM 1

Prove $f(n, k) = n + \binom{k}{2}$ for large n .

Alon (private communication): for large n , if the almost k -cover contains $x_1 = 1, \dots, x_n = 1$, then it contains at least $n + \binom{k}{2}$ affine hyperplanes in total.

n

$N_0 = [n]$

$(k-1)$ -subset

≥ 1

$N_0 \subseteq N_1 \subseteq N_0$

$\dots \in$

cover by hyperplane

PROBLEM 2

Let $g(n, m, k)$ be the minimum number of vertices covered less than k times by m affine hyperplanes not passing through $\vec{0}$. Determine $g(n, m, k)$.

Alon, Füredi 1993: $g(n, m, 1) = 2^{n-m}$.

Future research problems (II)

Question: Is it true that for all n, m, k :

$$g(n, m, k) = 2^{n-d},$$

where d is the maximum integer such that $f(d, k) \leq m$?

PROBLEM 3

Does there exist an absolute constant $C > 0$, which does not depend on n , such that for a fixed integer n , there exists M_n , so that whenever $k \geq M_n$,

$$f(n, k) \leq \left(1 + \frac{1}{2} + \cdots + \frac{1}{n}\right) k + C?$$

$$n=2, 3, 4$$



Thank you!